
Technical Information

Librel[®] Cu

A micronutrient fertilizer used to correct copper deficiency in crops and as a micronutrient source in growing media.

October 2021 | Supersedes issue dated August 2017 | Last change WF-No. 29259

10_100210e-03/Page 1 of 5

® = Registered trademark of BASF in many countries.


 **BASF**

We create chemistry


Product type	Micronutrient fertilizer conforming to the definition of an 'EC FERTILIZER'.
Description	Copper ethylenediamine tetra acetate disodium salt (CuEDTA Na ₂).
PRD-No.*	30482695 * BASF's commercial product number.

Intended use	To correct copper deficiency in most agricultural, horticultural and ornamental crops. Recommended for foliar or soil application. As a micronutrient source in hydroponics, liquid feed solutions and soilless growing media.
---------------------	--

Appearance	Librel® Cu is a blue spray agglomerated microgranule. The shade of blue will vary from batch to batch. Colour does not affect performance.
-------------------	--



Original Size



Enlarged Image

Handling and storage

Handling

- Librel® Cu should be stored indoors in a dry place.
- Care must be taken to exclude moisture. Drums/bags/boxes must be tightly resealed each time they are opened.
- Big bags are not to be stacked during storage in order to prevent lumping/agglomeration due to weight compression.
- Please refer to the latest Safety Data Sheet for detailed information on product safety.

Materials

Containers made of the following materials are appropriate for the storage of Librel® Cu:

- High density polyethylene (HDPE)
- Low density polyethylene (LDPE)
- Polypropylene (PP)

Shelf life

Librel® Cu has a shelf life of at least four years in its original packaging, provided it is stored correctly and drums/bags are kept tightly sealed.

Transport Precautions

No special precautions are necessary for transport by air, sea, rail or road.

Harmonised Tariff No.

2922 49 85 90


Properties

Some physical properties are listed in the table below. These are typical values only and not all of them are monitored on a regular basis. They are correct at the time of publication and do not necessarily form part of the product specification. A detailed product specification is available on request or via BASF's WorldAccount: <https://worldaccount.basf.com> (registered access).

Librel® Cu	Unit	Value
Physical form (25 °C)		microgranule
Cu (as water soluble Cu) (BASF method ICP-OES)	%	approx. 14.0
Dry weight BASF method, 110 °C)	%	approx. 95
Bulk density (BASF method)	g/l	approx. 900
Solubility (BASF method, in water @ 25 °C)	g/l	approx. 400
pH value (BASF method, 2% in water, 25 °C)		approx. 5.0

Particle size distribution

The following graph shows a typical particle size distribution curve for Librel® Cu, using Sympatec Helos H1594 Gradis System with an R7 lens:



X 10 µm	X 50 µm	VMD µm	X 90 µm	X 100 µm	Peak µm
126	229	244	377	730	235

Directions for use

General information

Librel® Cu gives best results when crops have adequate supplies of water and major nutrients and are not under stress for any other reason. Conditions which are responsible for one particular deficiency can also induce deficiencies of other micronutrients. Always ensure that deficiencies are confirmed before treatment is carried out.

Mixing with water

Simply add the powder to water while it is being agitated, do not pre-mix. Continue agitation for a short while to ensure complete dissolution.

Compatibility

Librel® Cu is compatible with all other Librel® chelates and many crop care chemicals. It is also fully compatible with solutions containing soluble phosphates such as liquid feeds and foliar fertilisers.

Foliar application

General information

Librel® Cu should be dissolved in a convenient volume of water to suit the spraying machine being used and the target crop leaf area. The following points should be observed.

1. The sprayer should be fitted with nozzles that produce a fine mist.
2. Only sufficient spray solution should be applied to coat the leaves and stems with a film of moisture with little or no “run off”.
3. Spraying should be carried out on a calm day **but not during strong sunshine or high temperatures**. The best time is late afternoon or evening.
4. If rain is imminent, spraying should be postponed. If rain falls within 4 hours of spraying, the crop should be re-sprayed 3 or 4 days later.

Fruit crops

Do not exceed a solution of 0.1% (1 g/l) for any one or combination of Librel® chelates. Some fruit varieties and cultivars can exhibit unpredictable sensitivity to EDTA chelates. Where local experience of successful use is not available, we strongly recommend small-scale test applications before wide spread use.

Rates of use

Crop	Rates of use (kg/ha)	Timing
Winter cereals	0.25 – 0.75	Apply as soon as active growth recommences in early spring. An additional half rate application at the flag stage (ZCK 37 – 39) is also beneficial
Other crops	0.25 – 0.75	Apply as soon as there is sufficient leaf area to absorb the spray

Water volume

The amount of Librel® Cu to be applied should be mixed with a volume of water appropriate to the crop leaf area of the type of spraying machine being used.

Arable crops: 200 – 600 litres per hectare.

NB: Do not exceed a solution concentration of 0.1% (1 g/l).

Wetting agent

Unless Librel® Cu is to be applied with a pesticide containing sufficient wetter, then a standard, agricultural non-ionic wetting agent should be used as recommended by the manufacturer.

Small scale use

For example using a knapsack sprayer. Prepare a 0.05 – 0.1% (0.5 – 1.0 g/l) solution and apply so as to coat the leaves and stems with a thin film of moisture with little or no run-off.

Soil application

General information

The appropriate amount of Librel® Cu should be dissolved in a convenient volume of water to suit the application equipment and to ensure even ground coverage.

General crops

Apply as a coarse low pressure spray immediately before the last cultivation prior to sowing or planting. Where crops are established, apply between the rows.

Perennial crops

Apply as a coarse low pressure spray in a wide circular band under the limit of the full branch spread. For best results, nutrients should be in the root zone before seasonal growth begins and this can be achieved by appropriate application timing.

Rates of use

Apply 0.5 – 2.0 kg/ha depending on the degree of deficiency crop size etc.

Hydroponic application

Rates of use

1 ppm (15.74 µmol/l) copper can be achieved by adding 7.15 g of Librel® Cu per 1000 litres of solution.

Statutory caution

To be used only where there is a recognised need. Do not exceed the appropriate dose rate.

Safety and Labelling

Please refer to the safety data sheet for information on classification & labelling, safe use, handling and transport.

Disclaimer

This document, or any answers or information provided herein by BASF, does not constitute a legally binding obligation of BASF. While the descriptions, designs, data and information contained herein are presented in good faith and believed to be accurate, it is provided for your guidance only. Because many factors may affect processing or application/use, we recommend that you make tests to determine the suitability of a product for your particular purpose prior to use. It does not relieve our customers from the obligation to perform a full inspection of the products upon delivery or any other obligation. NO WARRANTIES OF ANY KIND, EITHER EXPRESS OR IMPLIED, INCLUDING WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, ARE MADE REGARDING PRODUCTS DESCRIBED OR DESIGNS, DATA OR INFORMATION SET FORTH, OR THAT THE PRODUCTS, DESIGNS, DATA OR INFORMATION MAY BE USED WITHOUT INFRINGING THE INTELLECTUAL PROPERTY RIGHTS OF OTHERS. IN NO CASE SHALL THE DESCRIPTIONS, INFORMATION, DATA OR DESIGNS PROVIDED BE CONSIDERED A PART OF OUR TERMS AND CONDITIONS OF SALE.

Oktober 2021