



Cypro 200SC

For the control of diseases in turf, fruit and vegetable crops.

<u>Active Ingredient:</u>	200g/litre cyproconazole
<u>Chemical Group:</u>	Triazole/DMI – GROUP 3 FUNGICIDE
<u>Formulation:</u>	Suspension concentrate
<u>Pack Size:</u>	1 Litre

Cypro 200SC is a highly active DMI fungicide, with rapid activity that protects against a broad spectrum of diseases. It has long-lasting activity, ensuring the protection of new growth.



Cypro 200SC

– *The Proven DMI Performer:*

- Rapid penetration and long-lasting protection ✓
- Systemic and contact activity ✓
- Rapid rainfastness ✓
- Highly active, with very low use rates ✓

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

DIRECTIONS FOR USE:

Crop	Disease	Rate	Critical Comments
Grapes	Powdery Mildew (<i>Uncinula necator</i>)	7.5mL/100L	Apply non-DMI fungicides (eg sulphur) prior to flowering. From flowering apply Cypro 200SC at 2 to 3 week intervals. Ensure complete coverage using not less than 1000 litres water per hectare. Early season Cypro 200SC treatments may cause leaf discoloration on some grape varieties but this has not affected yield or fruit quality.
Peas	Powdery Mildew (<i>Erysiphe pisi</i>)	125mL/ha	Apply as a preventative spray when conditions favour disease development or at the first appearance of disease symptoms. If conditions persist for infection a repeat application may be necessary after 14 days.
Stonefruit	Brown Rot (blossom blight) (<i>Monilinia spp</i>)	7.5mL/100L	Make the first application at pink and repeat at 7 to 10 day intervals until shuck fall. Shorter intervals may be required in wet weather.
Vegetable and Field Brassicas	Ring Spot (<i>Mycosphaerella brassicicola</i>)	200mL/ha plus 2.5 L/ha 'all purpose' emulsifiable crop mineral oil	Apply as a preventative spray when conditions first favour disease development, or at the first sign of disease symptoms. Repeat at 2 to 3 week intervals. Apply a maximum of 4 applications per crop.

TURF GRASS DISEASE CONTROL			
Disease	Rate per 100m ²	Application Interval	Critical Comments
Brown Patch (<i>Rhizoctonia solani</i>)	15 – 20mL	21 – 28 days	Apply preventatively when conditions favour disease development, especially when warm temperatures and high humidity persist. When disease pressure is high use the shorter application interval and/or the higher rate. Repeat as necessary. DO NOT apply more than 4 applications of Cypro 200SC per year. Integrate with good turf management and a spray program with other fungicides. DO NOT apply more than 15 mL/100 square metres to creeping bent (<i>Agrostis stolonifera</i>) or <i>Poa annua</i> at intervals shorter than 28 days as phytotoxicity may occur. DO NOT apply to <i>Cotula</i> greens. Apply with properly calibrated equipment in 1000 to 2000 litres water/ha. DO NOT graze animals on treated turf areas, or use grass clippings for feed.
Dollar Spot (<i>Sclerotinia homeocarpa</i>)	10mL	21 – 28 days	
Rusts (<i>Puccinia spp</i>)	10mL	28 days	
Red Thread (<i>Laetisaria fuciformis</i>)	15 – 20mL	21 – 28 days	

MIXING: Add the required amount of Cypro 200SC to the half-filled spray tank. Add the remainder of the water and agitate thoroughly while mixing, spraying and when restarting after a stoppage.

COMPATIBILITY: Cypro 200SC is compatible with sulphur, mancozeb, streptomycin and most other fungicides and insecticides. Ensure thorough agitation of the mixture while spraying. DO NOT tank mix Cypro 200SC with copper fungicides.

IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE FOOD NOTICE: MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS.

Withholding Periods:

Grapes – 28 days

Peas – 14 days

Stonefruit – DO NOT apply after shuckfall

Brassicas – 14 days

Resistance Management: Cypro 200SC is a **GROUP 3 FUNGICIDE** from the DeMethylation (DMI/Triazole) group of fungicides. Resistance to Cypro 200SC and related DMI fungicides could develop from repeated use. To minimise this risk, use strictly in accordance with label instructions. **Grapes:** Use a maximum of 4 sprays, preferably in mixture with an approved protectant fungicide. No more than 2 DMI fungicides should be applied alone per season. **Stonefruit:** Use a maximum of 3 applications preferably in mixture with a non-DMI fungicide. Restrict use of the DMI fungicides to one part of the growing season, either during flowering or pre-harvest. Ensure full coverage of foliage.

Registered to and Distributed by: Adria New Zealand Limited. P.O. Box 535 Kumeu, Auckland 0841, NEW ZEALAND.

Ph: +64-9-412-9817; Fax: +64-9-412-9807. www.adria.nz

Cypro 200SC is Registered pursuant to the ACVM Act 1997, Approval No. P9401. EPA Approval Code: HSR101190.