

## TECHNICAL INFORMATION

### DuraTec® Top 14 14-7-14(+2+TE)



COMPO EXPERT GmbH  
Krögerweg 10  
48155 Münster  
Germany  
E-Mail: info@compo-expert.com

Technical data:	
Macronutrients	
<b>no EC - FERTILIZER</b>	
14,0 % nitrogen (N)	
8,0 % NH <sub>4</sub> nitrogen	
6,0 % NO <sub>3</sub> nitrogen	
- % NH <sub>2</sub> nitrogen	
- % isobutylienediurea	
- % crotonlyidenediurea	
- % methyleneurea	
7,0 % phosphate (P <sub>2</sub> O <sub>5</sub> ) soluble in neutral ammonium citrate and water	3,1 % P
5,6 % P <sub>2</sub> O <sub>5</sub> water soluble	2,4 % P
14,0 % potassium oxide (K <sub>2</sub> O), water soluble	11,6 % K
2,0 % magnesium (MgO)	1,2 % Mg
1,6 % water soluble MgO	1,0 % Mg
9,0 % sulphur (S)	22,5 % SO <sub>3</sub>
7,2 % water soluble S	18,0 % SO <sub>3</sub>
Micronutrients	
0,020 % boron (B)	- % copper (Cu)
0,060 % iron (Fe)	- % manganese (Mn)
0,010 % zinc (Zn)	- % molybdenum (Mo)
<b>low in chlorine</b>	

#### Other nutrients :

The raw materials used in the production process also contain low amounts of calcium (Ca), sodium (Na) or trace elements not mentioned above. The concentration of these elements is below EC declaration levels and can not be guaranteed.

#### Physical properties :

Colour :	purple + green
Bulk density :	1150 ± 100 kg / m <sup>3</sup>
Granule size :	90 % = 2 - 4 mm
pH ( 1:10 in water ) =	4,5 - 5,5
Physical appearance :	Granular solid fertilizer, surface-treated for improved transport and storage properties.

#### Recommendation for application :

Thanks to its even granulation the fertilizer can be applied easily with spreaders or by hand. Application is possible during the whole vegetation period. Watering-in promotes the immediate effect.

#### Description of fertilizer properties :

Controlled release NPK complex fertilizer with nitrification inhibitor DMPP (3,4 – dimethylpyrazolophosphate), partly coated.

The nutrients are released in two steps:

- 1) Starter effect after application under moisture influence.
- 2) In a second step, the nutrients are released out of the coated granules via diffusion influenced by moisture and temperature.

During the active phase of DMPP (4 to 10 weeks, depending on soil temperature and soil humidity) the transformation of ammonium to nitrate is delayed. As a result N-availability is further adapted to the plants` requirements and N-efficiency is increased.



For more information please contact your representative.

® registered trademark