

Recommendations for use

For stock fertilisation, PlantoSan® should be evenly mixed with the substrate before planting.

For top dressing, PlantoSan® should be evenly spread on bed and container surfaces.

The indicated rates of use apply to substrates containing no nutrients. If other fertilisers are to be added, the recommended rates of PlantoSan® should be reduced accordingly.

Vegetables - Seedling production		Top dressing of cut flowers	
Culture	Rates of use in kg/m³ or g/l of substrate	Cut flowers in the final place in original soil, e.g. Chrysanthemum, Cyclamen, Gerbera, Carnations, Roses : 40-60 g/m ²	
Celery	1.5		
Leek	1.5		
Cucumbers	1.5 - 2.0		
Head lettuce	1.5 - 2.0		
Peppers	1.5 - 2.0		
Savoy cabbage	1.5 - 2.0		
Cauliflower	2.0		
Brussels sprouts	2.0		
Cabbage red / white	2.0		
Tomatoes	2.0		
Seedling production of potted plants		Top dressing in tree nurseries	
Culture	Rates of use in kg/m³ or g/l of substrate	Fields of application	Rates of use g/l
<u>Plants with low nutrient requirement, e.g.:</u>		Container-grown plants	
Adiantum	0.75	- Top dressing in the 2nd year of cultivation	1 - 3
Anthurium scherzerianum hybr.	0.75	- Top dressing of plants in partly fertilised substrates	1 - 2
Primula accaulis	0.75		
Primula obconica	0.75		
<u>Plants with average nutrient requirement, e.g.:</u>		Fields of application	Rates of use g/m²
Anthurium andreaeanum hybr.	1.0	Surface fertilisation	
Begonia Elatior hybr.	1.0	- Shrubs (2 applications)	20 - 40
Cyclamen	1.0	- Broad-leaved trees and conifers (1-2 applications)	50
Euphorbia pulcherrima	1.0		
Sinningia (Gloxinia)	1.0		
<u>Plants with high nutrient requirement, e.g.:</u>		Top dressing in amenity areas	
Asparagus densiflorus	1.5	Fields of application	Rates of use* g/m²
Chrysanthemum	1.5	Ornamental shrubs in parks and public gardens	30 - 50
Codiaeum	1.5	Roses	30 - 50
Pelargonium	1.5	Roadside and dump plantation:	
		- 1-year-old seedlings	30
		- medium-sized bushes	50
		- bushes and stem bushes	50
		- half standards and standard trees	50
		*1 - 2 applications / year	
Production of bedding and balcony plants			
Culture	Rates of use in kg/m³ or g/l of substrate		
Pricking-out soils	1.0 - 2.0		
Seeding soils	0.5 - 1.0		

Distributor:



Horticulture - 0800 855 255
 TasmanCrop - 0800 855 255
 HortFertplus - 0800 273 748

Producer:



AGLUKON Spezialdünger GmbH & Co. KG
 Heerdtter Landstraße 199 · D-40549 Düsseldorf